DATE: 1217/91

SUPERFUND HEALTH RISK TECHNOLOGY SUPPORT CENTER at ECAO-Cincinnati

staffed by LABAT-ANDERSON, INC.

FAX NUMBER: (613) 569-7159 FTS: 684-7159

PHONE NUMBER: (613) 569-7300

FTS: 684-7300

TO: Mark Maddaloni	
LOCATION: Region II	
FAX NUMBER: FTS 264-6119	
COMMENTS:	
attached is the memo to John Osolin.	,
for the request submitted by arlene Lavin.	
a land copy will follow in the mail.	
O mindy Henson	
	_
	_
	_
280150	-
until nove ti-	_

DEC 17 191 14:19 STSCATHINECHOLCTM



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY OFFICE OF RESEARCH AND DEVELOPMENT ENVIRONMENTAL CRITERIA AND ASSESSMENT OFFICE CINCINNATI, OHIO 45268

DEC | 7 | 1991

SUBJECT:

Oral Toxicity Adjustment Factors and Dermal Absorption

Factors (Witco Chemical Site/Oakland, NJ)

FROM:

Kenneth A. Poirier

Director

Superfund Health Risk Technical Support Center

Chemical Mixtures Assessment Branch

TO:

John Osolin U.S. EPA Region II

This memorandum is in response to a request submitted by Arlene Levin of Alliance Technologies Corporation, for oral toxicity adjustment factors and dermal absorption factors for chemicals found at the Witco Chemical Superfund site.

Attached please find the following information:

Attachment I: Oral Adjustment Factors

Attachment II: Dermal Absorption Information

The absorption values provided in the attachments were obtained from available EPA documents and ATSDR Toxicological Profiles. References have been provided following the presentation of the tables.

There were a number of chemicals for which no information was available. Due to time constraints, we were unable to conduct a literature search for each one of those chemicals. Please notify us if you would like us to conduct additional research. This additional research would take us about 8 weeks to complete.

For chemicals listed as <u>ATSDR not available</u>, please contact ATSDR directly to obtain the most current ATSDR documents.

Please Note: No absorption information was provided in the attachments for lead. ECAO recommends you consult your EPA regional contact for appropriate regional guidance on assessing exposure to lead.

If you need further assistance, please feel free to contact the Superfund Technical Support Center at (513)569-7300 or FTS 684-7300.

Attachments

- J. Dinan (OS-230)

 - P. Grevatt (Region II)
 A. Levin (Alliance Technologies Corporation)
 M. Maddaloni (Region II)
 B. Means (OS-230)

Chemical

EPA Document

ATSDR Profiles

VOLATILES

Acetone (67-64-1)

Updated HEA 1989:
Animals: rats-by
gavage, acetone in
water rapidly
absorbed. HEA
1988: No
quantitative data;
readily absorbed

No ATSDR

Chloroform (67-66-5)

Updated HEA 1988: Humans: nearly complete. Mice, rats and Monkeys: 93-98%. HAD 1985: Animals: virtually completely absorbed. Rats, mice, monkeys 93-98% recovery of radioactivity in exhaled air, urine and carcass. AWQCD 1980 Animals: Approximately 100%

1991 Public Comment Draft: Humans: 100% absorbed rapidly. Animals: 93-98% absorption.

Chemical	cal <u>EPA Document</u>	
1,2,4-Trichloro- benzene (120-82-1)	HEA 1987 Animals: Rats and monkeys, 89-90% absorbed.	No ATSDR
Trichlorofluoro- methane (75-69-4)	DWHA 1989: No quantitative data, but expected to be absorbed.	No ATSDR
1,2,4-Trimethyl- benzene (95-63-6)	DWHA 1987: No quantitative data, but expected to be absorbed.	No ATSDR
1,3,5-Trimethyl- benzene	DWHA 1987: No quantitative data, but expected to be absorbed.	NO ATSDR
SEMIVOLATILES		
Benzo(a) anthracene	DWCD 1990: No data.	1990: Humans: No data. Animals: Rats: Rapid and efficient absorption.
Benzo(a)pyrene (50-32-8)	DWCD 1990: Rats: Approx. 60% in food; approx. 40% in starch suspension. HEA 1984: Rats: Approx. 50% from diet and starch suspension.	1990 Humans: No data. Rats: 1-3 % in urine and 74-85% in feces.

	Oral Absorption Factors		
<u>Chemical</u>	EPA Document	ATSDR Profiles	
Benzo(b)fluor- anthene (205-99-2)	DWCD 1990: No data.	1990: No quantitative data.	
Benzo(k)fluor- anthene (207-08-9)	No EPA Documents	1990: No quantitative data.	
Bis(2-ethyl hexyl)phthalate (117-81-7)	DWCD 1986: Animals: >90% from feed in rats. HEA 1987: Animals: >90% from feed in rats. Humans: 10-15% excreted in the urine as metabolites.	ATSDR 1989: Humans: minimum of 15%. Animals: >90% in rats.	
Butylbenzyl- phthalate (No cas#)	HEED 1989: Animals: Rats: 61-74% excreted in the urine. In ethanol/Emulphor 16.6% excreted in the urine. DWCD 1986: No quantitative data, on systemic effects following ingestion.	No ATSDR	

Chemical

EPA Document

ATSDR Profiles

Chrysene (218-01-9)

DWCD 1990:
(Polycyclic
Aromatic
Hydrocarbons):
Animals: 5-50% in
diet or as
suspension in
starch solution.
HEEP 1984:
Animals: Rats:
62.4-74.6%.

1990: Humans: No data. Animals: Absorbed but not quantified. Extent of absorption depends on the oral dose of chrysene and vehicle of administration. Approx. 25-41% was recovered from feces within 72 hrs. Olive oil -79% in feces following dietary and gavage administration.

Di-n-butylphthalate (No cas #)

HEEP 1987:
Animal: Rat >90%
absorption. HEA
1987: Animals:
Rats-gavage >90%
excreted as
metabolites in the
urine, the
remainder was
excreted in the
feces
DWCD 1986:
Animal: Rat >90%
absorption.

1991: Animal: Rat 63-97% absorption.

Fluoranthene (206-44-0)

DWCD 1990
(Polycyclic
Aromatic
Hydrocarbon): No
quantitative data.

1990: No quantitative data available.

Chemical

EPA Document

ATSDR Profiles

4-Nitrophenol (No cas #)

HEA 1987: (Nitrophenol): No pertinent quantitative data located in available literature. HEEP 1985: Pertinent data regarding absorption of 2nitrophenol, 3nitrophenol or 4nitrophenol were not located in available literature. AWOCD 1980: (Nitrophenols) No pertinent data.

ATSDR not available.

Phenanthrene (85-01-8)

DWCD 1990 (Polycyclic Aromatic Hydrocarbon): No quantitative data. HEA 1984: No specific data regarding GI absorption of phenanthrene are available. (Data from other structurally related PAHs suggest phenanthrene is readily absorbed / from GI tract) HEEP 1987: No pertinent data located.

1990: No quantitative data available.

Chemical

EPA Document

ATSDR Profiles

Phenol (108-95-2)

AWQCD 1980:
Readily absorbed
but not
quantified.
Updated HEA 1989:
Readily absorbed
but not
quantified.

1989: Humans: ~90% absorbed. Animals: Rats 95%; squirrel & monkey 31%.

Pyrene (129-00-0)

DWCD 1990
(Polycyclic
Aromatic
Hydrocarbon): No
quantitative data.
HEA 1984: Rats:
aqueous suspension
of pyrene was
poorly absorbed.
HEEP 1987:
Animals: Ratsabsorbed but not
quantified.

1989: No quantitative data available.

PESTICIDES/PCBs

4,4-DDD (72-54-8)

4,4-DDE (72-55-9) No relevant EPA documents.

AWQCD 1989: No data.
AWQCD 1980:
Absorbed with high efficiency. No quantitative data.

1990 Human:
Absorbed but not quantified.
Animal: 70-90 %.

Chemical EPA Document ATSDR Profiles 4,4-DDT HEA 1988: 1990: Human: Absorbed (50-29-3)Animals: rats estimated at 95% but not quantified. of absorption of Animal: 70-90%. dietary fats. Humans: slow but complete absorption. AWQCD 1989: No data. AWQCD 1980: 795% in dietary fats. Dieldrin 1991 Public HEA 1987: Humans: No data. Animal: Comment Draft: (60-57-1)Some absorption Humans: Doserelated response but not quantifiable. but not AWQCD 1980: quantifiable. (Aldrin/Dieldrin) Animal: Slow ³Cl-dieldrin: absorption but not Animals: Ratsquantifiable. immediately absorbed by upper GI tract. Absorption varies with solvent used. INORGANICS Aluminum ATSDR not HEA 1987: Humans: (7429 - 90 - 5)Absorbed but not available. quantified. Animals: Absorbed but not

quantified.

Chemical

EPA Document

ATSDR Profiles

Antimony (7440-36-0) AWQCD 1989: No quantitative data. Hea 1987: Human: Little absorption from GI tract (probably < 1%) for insoluble oxides. Animal: 15% absorption for water soluble organics.

1990 Public Comment Draft: Humans: 10% antimony tartrate. 1% all other forms.

Chemical

EPA Document

ATSDR Profiles

Arsenic (7440-38-2)

HEA 1984: Humans: >95% inorganic. Rats, pigs and monkeys: approx. 90% inorganic. AWQCD 1980: Must consider chemical form and its physical characteristics. Inorganic soluble arsenicsubstantial absorption of trivalent (>95%). Animal: either III or IV form is almost completely absorbed. Insoluble arsenictriselenide: poorly absorbed. Arsenic trioxide solution: Rabbits/Rats (in suspension): 40 & 30% abs. Monkeys: 98%. Pigs: 88%. Arsenate solution 70 to 90% abs. Shrimp or Seafood arsenic: Rapidly absorbed and rapidly excreted.

1989: Humans:
>95% inorganic.
Animals: >90%
water soluble
salts; 30-40%
trioxide
suspensions.

Barium (7440-39-3)

DWCD 1985:
Animals: 3-11%
gavage. HEA 1984:
Pertinent data not
located in
available
literature.

1990 Public Comment Draft: Humans: <5%. Animals: Dogs 7%; Rats <22 days old: 63-84%. Rats > 22 days old: 7%.

<u>Chemical</u>	EPA Document	ATSDR Profiles
Beryllium	HAD 1987: Human:	ATSDR not
(7440-41-7)	No data. Animal: <1% absorbed. DWCD	available.
	1991: < 1% absorption. Guinea pigs-0.006%	•
	absorption.	
Chromium (7440-47-3)	HAD 1984: Humans: 0.4% Cr III; 10.6% Cr VI. Animals: <3% Cr III. Rats: 2.3% Cr VI. HEA 1984: Humans: 2.1% Cr VI. Animals: 3-6% Cr VI. HAD 1984: No	1989: Humans: 0.4% Cr III; 10.6% Cr VI. Animals: 1.4- 3.0% Cr III; 2.4% Cr VI.
	pertinent data. AWQCD 1980: Low absorption,	
	<1% depending on chemical form in which the element is ingested.	
Chromium, total	No EPA Documents	No ATSDR

Chemical

EPA Document

ATSDR Profiles

Cobalt (7440-48-4)

HEA 1991 Revised 1st Draft: Human: 18-97% depending on dose of cobalt compound given and nutritional status of subjects. 31% abs. in irondeficient subjects. 18-44% in non-deficient subjects. Animal: ~30% gavage cobalt chloride. (more soluble compounds of cobalt were more readily absorbed than less soluble compounds).

1990 Public Comment Draft: Humans: 18-97% depending on type of cobalt, dose and nutritional status. Animals: Rats: 30%.

Copper (7440-50-8)

HEA 1984: Humans: 15-97% absorption. DWCD 1987: Human 15-97% absorption. AWQCD 1980: Humans: 40 to 60% dietary copper absorbed. Animals: Absorbed but not quantified. 1991: Humans: 15-97% copper acetate.

Iron (7439-89-6) HEA 1984: Humans: Heme Iron 10-25%; non heme 1-10%.

Magnesium (7439-95-4) No EPA Documents

No ATSDR

Chemical

EPA Document

ATSDR Profiles

Manganese (7439-96-5)

DWCD 1986:
Humans: Approx.
3%. Animals: Rats
approx. 3%. DWCD
1991: Human:
Studies ranging
from: -2.0±4.9%
to 8.4±4.7.
Calcium may
inhibit absorption
by increasing
alkalinity.
Animals: 3-4%.

1990 Public Comment Draft; Humans 3-5%. Rats: 2.5-5.5%.

Mercury (7439-97-6)

DWCD 1987: Humans: 78 mercuric nitrate 95% methyl mercury. Animals: Rats: 1-2% mercuric chloride 7% mercuric chloride in milk; suckling mice-38%; species not specified 0.01% (metallic). HEA 1984: Humans: High rate (methyl mercury); 15% (mercuric nitrate) AWQCD 1980: Virtually complete abs. Absorption of metallic mercury in liquid form is believed to be very small (<0.01% in animals). Inorganic mercury: human <15%.

1989: Humans: 0.1% (metallic), 15% (mercuric nitrate), 795% (methylmercuric nitrate). Animals: suckling mice - 38%, adult mice - 1% (standard diet), 7% (milk diet); mice-80% (organic mercury).

Chemical	EPA Document	ATSDR Profiles		
Nickel (7440-02-0)	HEA 1986: Humans: 1-10%. HAD 1985: Human: 1-10% absorption, dependent on dietary composition. AWQCD 1980: Human 1-10% absorption.	1988: Humans: 1-10%. Rats, dogs and mice: 1-10%		
Selenium (7782-49-2)	HEED 1989: Selenites, selenates and seleniferous compounds in mammals is rapid and efficient, but metal selenides & elemental selenium are absorbed poorly. HEA 1984: By gavage rats-91- 93% seleno- methionine. Humans: 70, 64, 44% selenite selenium. AWQCD 1980: Human: 70, 64, 44% of administered dose. Animal: Rats- selenite 91-93%. selenomethionine 95-97%.	1989: Humans: 80-97%. Animals: 80-100%.		
Silver (7440-22-4)	AWQCD 1989: Animals: 1-3% absorption.	ATSDR not available.		

Chemical

EPA Document

ATSDR Profiles

Thallium (7440-28-0)

AWQCD 1989 addendum: No data. AWQCD 1980: Completely absorbed, not quantified. ATSDR not available.

Vanadium (7440-62-2) HEA 1987: Humans: 0.1-1%. WQHA 1989: Human: 0.1-1.0% absorption with a mean of 0.5% rats fed vanadium sulfate.

1990 Public Comment Draft: Humans: No data. Animals: Rats 2.6%.

Zinc (7440-66-6) HEA 1984: (Zinc and compounds):
Human: 20-80% zinc chloride. Animal: very small amounts absorbed but not quantified.

ATSDR not available.

Chemical

EPA Documents

ATSDR Profiles

ATTACHMENT II

Volatiles

Acetone

Updated HEA 1989:

No information

available.

Chloroform

Updated HEA 1988: No information. HAD 1985: Slow absorption, no quantitative

1991 Public Comment Draft:

No ATSDR

Human: No data. Animal: No pertinent data.

information.

1,2-Dichlorobenzene

HEA 1987: No information.

No ATSDR

1,3-Dichlorobenzene HEA 1987: No information.

No ATSDR

1,4-Dichlorobenzene

HEA 1987: No information. AWQCD 1980:

Absorbed but not

quantified.

1991 Public

Comment Draft: No studies located for human or

animal.

Tetrachloroethylene

AWQCD 1980:

Human: Absorbed

but not quantified.

1991 Public Comment Draft:

Human: ≤1% absorption.

Animal: Absorbed

but not specifically quantified.

1,2,4-Trichlorobenzene

HEA 1987: No information.

No ATSDR

Trichlorofluormethane

No EPA Documents

No ATSDR

Chemical

EPA Documents

ATSDR Profiles

1,2,4-Trimethylbenzene DWHA 1987: Percutaneous absorption but not quantified. No ATSDR

1,3,5-Trimethylbenzene DWHA 1987: Percutaneous absorption but not quantified. No ATSDR

SEMIVOLATILES

Benzo(a) anthracene

(PAHS) DWCD 1990: Absorption dependent on molecular size, structure and the nature of the solvent. Small amounts of mineral oil enhance dermal absorption. However, the absorption was not as great when a mixture of 50% mineral oil was used as the solvent.

1990 Human: information. Animal: Rats-52.3% absorption. Over time, permeation significantly decreased suggesting it is dermally absorbed in a dosedependent manner. Diffusion through skin depended on the amount of anthracene on skin's surface.

Chemical

EPA Documents

ATSDR Profiles

Benzo(a) pyrene

DWCD 1990: (PAHs)
Animal: Rats 90%
absorption.
Depends on size
and configuration
of the hydrocarbon
molecule and the
concentration of
hydrocarbon
applied to the
skin.

1990: An in vitro study using human skin revealed 3% absorption after 24hrs. Animal: 6% and 40%, at 1 and 24 hours following administration, respectively. Seven days after exposure 93% was recovered in the feces.

Benzo(b) fluoranthene DWCD 1990: (PAHs) No information.

Benzo(k) fluoranthene DWCD 1990: (PAHs) No information.

Bis(2-ethylhexyl)phthalate No EPA Documents

No ATSDR

Butylbenzylphthalate HEED 1989: No information.

No ATSDR

Chrysene

HEEP 1984: No information. DWCD 1990: No information.

1990: Human: No information. Animal: Absorbed but not quantified.

Di-n-butylphthalate DWCD 1991: (Phthalic Acid Esters) No information.

ATSDR not available.

Fluoranthene

DWCD 1990: No information.

1990: No information.

ATSDR not

1990: No

information.

available.

	em	
~		

EPA Documents

ATSDR Profiles

4-Nitrophenol

HEA 1987: (Nitrophenols) -No information. HEEP 1985:

(Nitrophenols) -Animals: Dogs: ~11% urine; <0.5% feces. Rabbits~35%

urine; 0.53%

feces.

AWQCD 1980: No pertinent

information.

Phenanthrene

HEEP 1987: Highly soluble can pass across epithelial membranes-no quantitative data.

HEA 1984: No information.

Phenol

AWQCD 1980: Readily absorbed but not

quantified.

Pyrene

HEEP 1987: No information. HEA 1984: No information.

1990: No information.

ATSDR not

available.

Pesticides/PCBs

4,4-DDD

No relevant EPA documents.

ATSDR not available.

4,4-DDE

AWQCD 1989: No information. AWQCD

1980: No information.

ATSDR not available.

Chemical	EPA Documents	ATSDR Profiles		
4,4-DDT	AWQCD 1989:No information. AWQCD 1980: No information.	ATSDR not available.		
Dieldrin	AWQCD 1980: No information. HEA 1987: No information.	1991 Public Comment Draft: Humans: Rapid absorption-7.7% but low confidence in study cited. Animals: Rats: Rapidly absorbed but not quantified.		
INORGANICS		•		
Aluminum	HEA 1987: No information.	ATSDR not available.		
Antimony	HEA 1987: No information.	ATSDR not available.		
Arsenic	AWQCD 1989: No information. AWQCD 1980: Human: No pertinent data. Animal: Rats- solutions of arsenate, significant absorption but not	1991 Public Comment Draft: No quantitative studies were located on inorganic arsenic.		
Barium	quantified. HEA 1984: No	ATSDR not		
 	information.	available.		

Chem	34	<u>.</u>	•	7
CIICH			-	<u> </u>

EPA Documents

ATSDR Profiles

marina a dame	DWCD 1991:	ATSDR not
Beryllium	Animals: Rats-	available.
	poorly absorbed	
•	although contact	•
		•
	dermatitis can	•
	occur.	
• •	HEA 1987: No	**
	information.	
	HEA 1984: No	
	information.	
	AWQCD 1980: No	
	information.	
Chromium (VI)	HEA 1984: No	ATSDR not
	information.	available.
•	HAD 1984:	
	Absorbed but not	
	quantified.	
	AWQCD 1980:	
	(Chromium) VI	
	fairly readily	
	absorbed however	
•	not quantified.	, · ·
• •		• ',
Chromium (Total)		ATSDR not
	. '	available
Cobalt	HEA 1991 Revised	ATSDR not
OODGIC	1st Draft: No	available.
	data.	
	•	
Copper	DWCD 1987: No	ATSDR not
	pertinent	available.
•	information.	
	AWQCD 1980: No	
•	pertinent data.	
	has arreste ander	
Iron	HEA 1984: No	No ATSDR
*** ***	information.	
	THE OF THE PERIOD	,
Magnesium	No EPA documents.	No ATSDR
	TA STATE MANAGEMENT	

Chem	i	c	à	1

EPA Documents

ATSDR Profiles

ATSDR not DWCD 1991: No Manganese available. information. ATSDR not AWQCD 1989: No Mercury available. information. AWQCD 1980: Inorganic mercury: No quantitative information. Mercuric chloride: Animals: 5%. Humans: expect substantial absorption but not quantified. AWQCD 1980: Nickel Human: absorbed but not quantified.

Selenium AWQCD 1989: No information.
HEED 1989: No information.
HEA 1984: No information.
AWQCD 1980: No pertinent information -

absorption enhanced by sweat and detergents.

Thallium AWQCD 1989: No information.

AWQCD 1980:

Absorbed but not quantified.

ATSDR not available.

ATSDR not

available.

Chemical

EPA Documents

ATSDR Profiles

Vanadium

WQHA 1989:

ATSDR not available.

Animals: Rabbits: absorbed but not quantified.

Human: absorbed but not quantified.

Zinc

HEA 1984: No information.

ATSDR not available.

References

HEA - Health Effects Assessment

- U.S. EPA. 1989. Updated Health Effects Assessment for Acetone. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1987. Health Effects Assessment for Aldrin/Dieldrin. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1987. Health Effects Assessment for Aluminum. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1987. Health Effects Assessment for Antimony.
 Prepared by the Office of Health and Environmental Assessment,
 Environmental Criteria and Assessment Office, Cincinnati, OH for
 the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1984. Health Effects Assessment for Arsenic. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1984. Health Effects Assessment for Barium. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1984. Health Effects Assessment for Benzo[a]pyrene. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
 - U.S. EPA. 1987. Health Effects Assessment for Beryllium. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
 - U.S. EPA. 1984. Health Effects Assessment for Beryllium. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
 - U.S. EPA. 1988. Updated Health Effects Assessment for Chloroform. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office,

- Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1984. Health Effects Assessment for Chromium. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1991. Revised First Draft. Health Effects Assessment for Cobalt. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1984. Health Effects Assessment for Copper. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1988. Health Effects Assessment for 4,4-DDT.
 Prepared by the Office of Health and Environmental Assessment,
 Environmental Criteria and Assessment Office, Cincinnati, OH for
 the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1987. Health Effects Assessment for Dichlorobenzenes. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1984. Health Effects Assessment for Iron. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1984. Health Effects Assessment for Mercury.
 Prepared by the Office of Health and Environmental Assessment,
 Environmental Criteria and Assessment Office, Cincinnati, OH for
 the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1986. Health Effects Assessment for Nickel. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1987. Health Effects Assessment for Nitrophenols. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1984. Health Effects Assessment for Phenanthrene. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for

- the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1987. Health Effects Assessment for Bis (2-ethylhexyl)Phthalate. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1987. Health Effects Assessment for Di-n-butyl Phthalate. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1984. Health Effects Assessment for Pyrene. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1984. Health Effects Assessment for Selenium.
 Prepared by the Office of Health and Environmental Assessment,
 Environmental Criteria and Assessment Office, Cincinnati, OH for
 the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1988. Health Effects Assessment for Tetrachloroethylene. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1987. Health Effects Assessment for 1,2,4-Trichlorobenzene. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1987. Health Effects Assessment for Vanadium. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1987. Health Effects Assessment for Zinc and Compounds. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Emergency and Remedial Response, Washington, DC.

Ambient Water Quality Criteria Document (AWOCD)

- U.S. EPA. 1980. Ambient Water Quality Criteria for Aldrin/Dieldrin. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1989 Addendum. Ambient Water Quality Criteria for Antimony. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1989 Addendum. Ambient Water Quality Criteria for Arsenic. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1980. Ambient Water Quality Criteria for Arsenic. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1980. Ambient Water Quality Criteria for Beryllium. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1980. Ambient Water Quality Criteria for Chloroform. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1980. Ambient Water Quality Criteria for Chromium. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1980. Ambient Water Quality Criteria for 4,4-DDE. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1989 Addendum. Ambient Water Quality Criteria for 4,4-DDE. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.

- U.S. EPA. 1980. Ambient Water Quality Criteria for 4,4-DDT. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1989 Addendum. Ambient Water Quality Criteria for 4,4-DDT. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1980. Ambient Water Quality Criteria for 1,4-Dichlorobenzene. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1989 Addendum. Ambient Water Quality Criteria for Mercury. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1980. Ambient Water Quality Criteria for Mercury. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1980. Ambient Water Quality Criteria for Nickel. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1980. Ambient Water Quality Criteria for Nitrophenols. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1980. Ambient Water Quality Criteria for phenol. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1989 Addendum. Ambient Water Quality Criteria for Selenium. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1980. Ambient Water Quality Criteria for Selenium. Prepared by the Office of Health and Environmental Assessment,

- Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1989 Addendum. Ambient Water Quality Criteria for Selenium. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1980. Ambient Water Quality Criteria for Tetrachloroethylene. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1980. Ambient Water Quality Criteria for Thallium. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1989 Addendum. Ambient Water Quality Criteria for Thallium. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.

Health Assessment Document (HAD)

- U.S. EPA. 1987. Health Assessment Document for Beryllium.
 Office of Health and Environmental Assessment, Environmental
 Criteria and Assessment Office, Cincinnati, OH for Office of Air
 Quality Planning and Standards, Washington, DC.
- U.S. EPA. 1985. Health Assessment Document for Chloroform. Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for Office of Air Quality Planning and Standards, Washington, DC.
- U.S. EPA. 1984. Health Assessment Document for Chromium.
 Office of Health and Environmental Assessment, Environmental
 Criteria and Assessment Office, Cincinnati, OH for Office of Air
 Quality Planning and Standards, Washington, DC.
- U.S. EPA. 1985. Health Assessment Document for 1,3,5-Dichlorobenzene. Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for Office of Air Quality Planning and Standards, Washington, DC.

- U.S. EPA. 1985. Health Assessment Document for Nickel. Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for Office of Air Quality Planning and Standards, Washington, DC.
- U.S. EPA. 1984. Health Assessment Document for Chromium.
 Office of Health and Environmental Assessment, Environmental
 Criteria and Assessment Office, Cincinnati, OH for Office of Air
 Quality Planning and Standards, Washington, DC.

Health and Environmental Effects Profile (HEEP)

- U.S. EPA. 1984. Health and Environmental Effects Profile for Chrysene. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH, for the Office of Solid Waste and Emergency Response, Washington, DC.
- U.S. EPA. 1985. Health and Environmental Effects Profile for Nitrophenols. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH, for the Office of Solid Waste and Emergency Response, Washington, DC.
- U.S. EPA. 1987. Health and Environmental Effects Profile for Phenanthrene. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH, for the Office of Solid Waste and Emergency Response, Washington, DC.
- U.S. EPA. 1987. Health and Environmental Effects Profile for Di-n-butyl Phthalate. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH, for the Office of Solid Waste and Emergency Response, Washington, DC.
- U.S. EPA. 1987. Health and Environmental Effects Profile for Pyrene. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH, for the Office of Solid Waste and Emergency Response, Washington, DC.
- U.S. EPA. 1989. Health and Environmental Effects Profile for Selenium. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH, for the Office of Solid Waste and Emergency Response, Washington, DC.

Health and Environmental Effects Document (HEED)

- U.S. EPA. 1989. Health and Environmental Effects Document for Butylbenzyl Phthalate. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH, for the Office of Solid Waste and Emergency Response, Washington, DC.
- U.S. EPA. 1989. Health and Environmental Effects Document for Selenium. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH, for the Office of Solid Waste and Emergency Response, Washington, DC.

Drinking Water Health Advisory (DWHA)

- U.S. EPA. 1989. Trichlorofluoromethane. Health Advisory. Office of Drinking Water, Washington, DC.
- U.S. EPA. 1987. 1,2,4-Trimethylbenzene. Health Advisory. Office of Drinking Water, Washington, DC.
- U.S. EPA. 1987. 1,3,5-Trimethylbenzene. Health Advisory. Office of Drinking Water, Washington, DC.

Water Quality Health Advisory (WOHA)

U.S. EPA. 1989. Water Quality Health Advisory for Vanadium. Prepared by The Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.

Drinking Water Criteria Document (DWCD)

- U.S. EPA. 1985. Drinking Water Criteria Document for Barium. Prepared by Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Drinking Water, Washington, DC.
- U.S. EPA. 1991. Drinking Water Criteria Document for Beryllium. Prepared by Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Drinking Water, Washington, DC.
- U.S. EPA. 1986. Drinking Water Criteria Document for Bis (2-ethylhexyl) Phthalate. Prepared by Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Drinking Water, Washington, DC.

- U.S. EPA. 1986. Drinking Water Criteria Document for Butylbenzyl Phthalate. Prepared by Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Drinking Water, Washington, DC.
- U.S. EPA. 1987. Drinking Water Criteria Document for Copper. Prepared by Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Drinking Water, Washington, DC.
- U.S. EPA. 1990. Drinking Water Criteria Document for Fluoranthene. Prepared by Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Drinking Water, Washington, DC.
- U.S. EPA. 1991. Drinking Water Criteria Document for Manganese. Prepared by Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Drinking Water, Washington, DC.
- U.S. EPA. 1986. Drinking Water Criteria Document for Manganese. Prepared by Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Drinking Water, Washington, DC.
- U.S. EPA. 1987. Drinking Water Criteria Document for Mercury. Prepared by Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Drinking Water, Washington, DC.
- U.S. EPA. 1990. Drinking Water Criteria Document for PAHs. Prepared by Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Drinking Water, Washington, DC.
- U.S. EPA. 1986. Drinking Water Criteria Document for Phthalic Acid Esters. Prepared by Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Drinking Water, Washington, DC.
- U.S. EPA. 1991. Drinking Water Criteria Document for Phthalic Acid Esters. Prepared by Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Drinking Water, Washington, DC.